

Tables

Table 5.1. Area occupied by sagebrush in the Intermountain Region estimated from Küchler's map of potential vegetation (Küchler 1970 and West 1983). The regional boundaries used for these area estimates were delineated in Küchler's map (1970) of potential vegetation (Fig. 5.3) and did not include eastern parts of sagebrush biome within the Conservation Assessment area.

| State | Area (km ²) | Percentage of total area |
|-----------------------|-------------------------|-----------------------------|
| Sagebrush Steppe | | |
| Wyoming | 109,000 | 24.3 |
| Idaho | 103,000 | 22.9 |
| Oregon | 92,000 | 20.5 |
| Nevada | 47,000 | 10.6 |
| Washington | 38,000 | 8.5 |
| California | 18,000 | 4.1 |
| Colorado | 17,000 | 3.8 |
| Montana | 13,000 | 2.9 |
| Utah | 11,000 | 2.4 |
| Total | 448,000 | 100.0 |
| Great Basin Sagebrush | | |
| Nevada | 106,000 | 59.2 |
| Utah | 27,000 | 15.1 |
| Arizona | 24,000 | 13.5 |
| Colorado | 9,000 | 5.1 |
| California | 8,000 | 4.4 |
| New Mexico | 5,000 | 2.7 |
| Total | 179,000 | 100.0 |